

Facts on Moles



Damage caused by Moles

Mole hills can spoil the look of a lawn, park or golf course. Stones can damage grass cutting blades and soil will spoil grass silage.

Territory

Each mole has its own burrow system, a network of firm-walled tunnels. The tunnels are about 5cm wide, 4cm high and may be over 70 metres long; they vary in depth from just beneath the surface of the ground to about 70cm. When digging close to the surface, the moles push the displaced soil up vertical tunnels and these form the familiar molehills. The territories of several moles may overlap, but the residents avoid each other if they can, except in the breeding season. If two males meet, they may fight fiercely, which can result in death.

Daily Life

Moles are active by day and by night, almost continuously digging their tunnels and searching for food. They are active for about four hours at a time and then rest for a similar length of time. A mole will die of starvation if it does not eat every few hours.

It finds food by running along its tunnels and eating up any earthworms, beetle larvae, slugs etc, which have fallen from the walls. A mole probably eats at least half its own body weight a day. When earthworms are plentiful, during autumn and winter, the mole makes stores of them, biting off their heads and pushing

them into the ground for eating later.

A mole can run backwards through tunnels and turn right round by doing a somersault! Its velvety fur lies backwards or forwards so that it does not become stuck against the tunnel walls when squeezing through them.

The eyes of the mole are very tiny but sight is not important to an animal that lives in darkness for most of the time. It does not have a good sense of smell or hearing but it is extremely sensitive to touch and can sense vibrations in the soil around it. Sensitive whiskers help it to find its way about and detect food and water; it is also helped by thousands of very sensitive, tiny hairs which cover its long nose.

Moles are sometimes seen above ground. They come to the surface to collect nesting material and to look for food when the soil is dry. Young moles come to the surface to look for new homes when they leave their mother's burrow. Moles emerge mainly at night but they are still vulnerable to predators; some mammals find them distasteful but many are eaten by tawny and barn owls.

Breeding

Moles breed from March to May only. The male (boar) visits the female (sow) in her burrow and leaves straight after mating, taking no part in raising the young. The female builds a special chamber about the size of a football and lines it with dry grass and leaves. An extra large molehill, or 'fortress', usually covers the nest chamber. A food store will be close by.

After a gestation period (time between mating and birth) of 30 days, a litter of 2-7 young is born. The babies are blind and naked at birth and do not grow fur until they are 2 weeks old. At about 3 weeks of age their eyes open. They grow quickly, feeding on their mother's milk, and are ready to leave the nest at about 35 days. They travel overland searching for territories of their own.

For more detailed information and a free telephone consultation call 0845 838 0634.

Specialists in Pest and Vermin Control